

Isomorphism classes of Galois extensions

Robert G. Underwood
Department of Mathematics
Department of Computer Science
Auburn University at Montgomery
Montgomery, Alabama



May 28, 2026

Contents

1. Galois extensions
2. The Induction map of Pareigis
3. The Haggemüller-Pareigis bijection
4. Hopf-Galois Structures
5. Another Example

1. Galois extensions

Throughout this talk, K is a finite extension of \mathbb{Q} .

The notion of a Galois extension of K is due to M. Auslander and O. Goldman [AG60].

Let A be a finite dimensional commutative K -algebra and let $\text{End}_K(A)$ denote the K -algebra of K -linear maps $\phi : A \rightarrow A$. Let $\text{Aut}_K(A)$ denote the group of K -algebra automorphisms of A .

Let F be a finite subgroup of $\text{Aut}_K(A)$ with $K = A^F$. Let $D(A, F)$ denote the crossed product algebra of A by F .

Then A is an F -Galois extension of K if the map

$$j : D(A, F) \rightarrow \text{End}_K(A),$$

defined as $j(\sum_{g \in F} a_g g)(t) = \sum_{g \in F} a_g g(t)$, $a_g, t \in A$, is an isomorphism of K -algebras.

The definition of F -Galois extension generalizes the usual definition of a Galois extension of fields.

Let A, A' be F -Galois extensions of K . Then A is isomorphic to A' as F -Galois extensions of K if there exists an isomorphism of commutative K -algebras $\theta : A \rightarrow A'$ for which

$$\theta(g(x)) = g(\theta(x))$$

for all $g \in F, x \in A$.

Let $\mathcal{Gal}(K, F)$ denote the set of isomorphism classes of F -Galois extensions of K .

Example 1.

Let $\text{Map}(F, K)$ denote the K -algebra of maps $\phi : F \rightarrow K$. Then $\text{Map}(F, K)$ is the *trivial F -Galois extension* of K with action defined as

$$g(\phi)(h) = \phi(g^{-1}h)$$

for $g, h \in F$, $\phi \in \text{Map}(F, K)$.

The set of maps $\{v_g\}_{g \in F}$ defined as

$$v_g(h) = \delta_{g,h},$$

for $g, h \in F$, is a K -basis for $\text{Map}(F, K)$.



The Galois extensions are completely determined by the following.

Theorem 2.

Let K be a field, let F be a finite group and let A be an F -Galois extension of K . Then

$$A = \underbrace{L \times L \times \cdots \times L}_n$$

where L is a U -Galois field extension of K for some subgroup U of F of index n . (L is a Galois extension of K with group U in the usual sense.)

Proof. See[Pa90, Theorem 4.2].



One of the main points of this talk is to answer the following question: Can we recover A from the data U, L ? In other words: can A be induced from the data U, L ?

The answer is yes (of course), and to do this we need the induction map of Pareigis.

2. The Induction map of Pareigis

Let $U \leq F$ and suppose that $M \in \mathcal{G}al(K, U)$ with $n = [F : U]$.

Let $\{g_1, g_2, \dots, g_n\}$ be a left transversal for U in F and let $\{g_1 U, g_2 U, \dots, g_n U\}$ be the set of left cosets of U in F . We take $g_1 = 1$. Let

$$A = \underbrace{M \times M \times \dots \times M}_n$$

with minimal orthogonal idempotents $\epsilon_1, \epsilon_2, \dots, \epsilon_n$.

Let $\varsigma : F \rightarrow S_n$ be defined as $\varsigma(f)(i) = j$ if $fg_i U = g_j U$.

Define an action of F on A as follows: for $f \in F$,

$$f\left(\sum_{i=1}^n m_i \epsilon_i\right) = \sum_{i=1}^n (g_{\sigma(f)(i)}^{-1} f g_i)(m_i) \epsilon_{\sigma(f)(i)},$$

for $m_i \in M$, $1 \leq i \leq n$.

Then A is an F -Galois extension of K [Pa90, Theorem 4.2].

In this way, we define a map

$$A(U, \) : \mathcal{G}al(K, U) \rightarrow \mathcal{G}al(K, F),$$

$M \mapsto \underbrace{M \times M \times \cdots \times M}_n$, which is the *Pareigis induction map*.

Theorem 3.

Let F be a finite group. Let A be an F -Galois extension of K . Then there exists a subgroup $U \leq F$ and a U -Galois extension L of K for which

$$A \cong A(U, L/K),$$

as F -Galois extensions of K , i.e., A can be induced from the data $U, L/K$.

Proof. By Theorem 2,

$$A = \underbrace{L \times L \times \cdots \times L}_n, \quad (1)$$

where L is a U -Galois field extension of K for some subgroup U of F of index n .

Now with the data $U, L/K$, we construct an F -Galois extension $A(U, L/K)$ using the induction map of Pareigis:

$$A(U, L/K) = \underbrace{L \times L \times \cdots \times L}_n.$$

We then show that $A \cong A(U, L/K)$ as F -Galois extensions of K .

□

Proposition 4.

Let $U, V \leq F$ be distinct subgroups that are isomorphic. Let L/K be a Galois field extension with both U and V as its Galois group. Then $A(U, L/K) \cong A(V, L/K)$ as F -Galois extensions of K if and only if U, V are conjugate.

Proof. Suppose that U, V are conjugate. Then there exists $f \in F$ with $U = fVf^{-1}$. Set $h_1 = f$ and let $\{fV, h_2V, h_3V, \dots, h_nV\}$ denote the set of distinct left cosets of V in F . The group F acts transitively on $\{fV, h_2V, h_3V, \dots, h_nV\}$.

For $u \in U$, $u = fvf^{-1}$ for some $v \in V$, thus

$$ufV = fvf^{-1}fV = fvV = fV.$$

Moreover, if $gfV = fV$, then for $v \in V$, $gfv = fv'$ for some $v' \in V$. So, $gf = fv''$, $v'' \in V$, so $g = fv''f^{-1} \in U$.

Thus U stabilizes fV ; fV is the “base point.”

Consequently, there is an isomorphism of F -sets

$$\alpha : \{fV, h_2V, \dots, h_nV\} \rightarrow \{g_1U, g_2U, \dots, g_nU\},$$

defined as $\alpha(h_iV) = g_iU$, where $g_i \in F$ satisfies $g_ifV = h_iV$.

Next, we define a map

$$\theta : A(V, L/K) \rightarrow A(U, L/K)$$

by the rule

$$\theta\left(\sum_{i=1}^n l_i \epsilon_i\right) = \sum_{i=1}^n g_i^{-1}(l_i) \alpha(\epsilon_i),$$

for $l_i \in L/K$, $1 \leq i \leq n$.

Now, for $f \in F$,

$$\begin{aligned}
f(\theta(\sum_{i=1}^n l_i \epsilon_i)) &= f(\sum_{i=1}^n g_i^{-1}(l_i) \alpha(\epsilon_i)) \\
&= \sum_{i=1}^n (g_i^{-1} f g_i)(g_i^{-1}(l_i)) \alpha(\epsilon_i) \\
&= \sum_{i=1}^n g_i^{-1}(f(l_i)) \alpha(\epsilon_i) \\
&= \theta(\sum_{i=1}^n f(l_i) \epsilon_i) \\
&= \theta(f(\sum_{i=1}^n l_i \epsilon_i)).
\end{aligned}$$

Thus $A(U, L/K) \cong A(V, L/K)$ as F -Galois extensions of K .

For the converse, assume that $A(U, L/K) \cong A(V, L/K)$ as F -Galois extensions of K . Then there exists an F -invariant isomorphism of F -sets

$$\alpha : \{U, g_2 U, \dots, g_n U\} \rightarrow \{V, h_2 V, \dots, h_n V\}.$$

We have $\alpha(U) = hV$ for some $h \in F$ and so, $uhV = hV$ for all $u \in U$. Let $u \in U, v \in V$. Then

$$h^{-1}uhv = v'$$

for some $v' \in V$.

Thus $h^{-1}uh = v'v^{-1} \in V$. If u, u' are distinct elements of U , then $h^{-1}uh$ and $h^{-1}u'h$ are distinct elements of V . Since $|U| = |V|$, we conclude that $h^{-1}Uh = V$, i.e., U, V are conjugate.

□

Example 5. Let $F = S_3$ with presentation

$$S_3 = \langle a, b : a^3 = b^2 = 1, ba = a^2b \rangle.$$

There are 3 isomorphic subgroups of S_3 of order 2:

$$U_1 = \{1, b\}, \quad U_2 = \{1, ba\}, \quad U_3 = \{1, ab\}.$$

Consider the Galois extension $\mathbb{Q}(\sqrt{2})$ of \mathbb{Q} , and consider each U_i , $1 \leq i \leq 3$, to be its Galois group.

Since $[F : U_i] = 3$ for $1 \leq i \leq 3$, each $A(U_i, \mathbb{Q}(\sqrt{2})/\mathbb{Q})$ appears as

$$\mathbb{Q}(\sqrt{2}) \times \mathbb{Q}(\sqrt{2}) \times \mathbb{Q}(\sqrt{2}),$$

and thus the $A(U_i, \mathbb{Q}(\sqrt{2})/\mathbb{Q})$ are pairwise isomorphic as \mathbb{Q} -algebras.

The subgroups U_i are pairwise conjugate. Thus by Proposition 4, $A(U_i, \mathbb{Q}(\sqrt{2})/\mathbb{Q})$ are pairwise isomorphic as S_3 -Galois extensions of \mathbb{Q} .



3. The Haggemüller-Pareigis bijection

Let N be a finite group with automorphism group $F = \text{Aut}(N)$.

Let $K[N]$ denote the group ring K -Hopf algebra.

Let $\mathcal{F}orm(K[N])$ denote the collection of the isomorphism classes of the forms of $K[N]$.

Theorem 6 (Haggenmüller and Pareigis).

There exists a bijection

$$\Theta : \mathcal{G}al(K, F) \rightarrow \mathcal{F}orm(K[N]),$$

which gives a 1-1 correspondence between the isomorphism classes of F -Galois extensions of K and forms of $K[N]$, where $F = \text{Aut}(N)$. For an F -Galois extension A of K , the map Θ is given explicitly as the fixed ring

$$\Theta(A) = (A[N])^F,$$

where F acts on A through the Galois action and on N as automorphisms. The fixed ring $(A[N])^F$ is an A -form of $K[N]$ and so belongs to $\mathcal{F}orm(K[N])$.

Proof. See [HP86, Theorem 5].



Example 7. Let $N = C_2 \times C_2$. Then $F = \text{Aut}(C_2 \times C_2) = S_3$.

In Example 5, we constructed a collection of S_3 -Galois extensions of \mathbb{Q} :

$$A(U_i, \mathbb{Q}(\sqrt{2})/\mathbb{Q}),$$

$1 \leq i \leq 3$. The $A(U_i, \mathbb{Q}(\sqrt{2})/\mathbb{Q})$ are pairwise isomorphic as S_3 -Galois extensions.

Let

$$\Theta : \mathcal{G}al(\mathbb{Q}, S_3) \rightarrow \mathcal{F}orm(\mathbb{Q}[C_2 \times C_2])$$

be the Hagenmüller-Pareigis bijection.

Now, the set of images

$$\Theta(A(U_i, \mathbb{Q}(\sqrt{2})/\mathbb{Q})) = A(U_i, \mathbb{Q}(\sqrt{2})/\mathbb{Q})[C_2 \times C_2]^{S_3}$$

are pairwise isomorphic as \mathbb{Q} -Hopf algebras (and as Hopf forms of $\mathbb{Q}[C_2 \times C_2]$).



4. Hopf-Galois structures

By Greither/Pareigis, certain Hopf forms of $K[N]$ arise as Hopf-Galois structures on Galois field extensions.

Let E/K be a Galois extension with group G , and let $\lambda : G \rightarrow \text{Perm}(G)$ denote the left regular representation.

Let (H_N, \cdot_N) be a Hopf-Galois structure on E/K corresponding to a regular subgroup $N \leq \text{Perm}(G)$ that is normalized by $\lambda(G)$.

Necessarily, $|N| = |G|$, yet we may have $N \not\cong G$ as groups; the isomorphism class of N is the *type* of (H_N, \cdot_N) .

Let $(H_{N'}, \cdot_{N'})$ be some other Hopf-Galois structure on E/K corresponding to a regular subgroup $N' \leq \text{Perm}(G)$ normalized by $\lambda(G)$. Then $(H_{N'}, \cdot_{N'})$ is of the *same type* as (H_N, \cdot_N) if $N' \cong N$.

The K -Hopf algebra H_N of the Hopf-Galois structure is a Hopf-algebra E -form of $K[N]$, which by Galois descent corresponds to a homomorphism

$$\varrho_N : G \rightarrow F_N$$

in

$$H^1(G, \mathbf{Aut}(K[N])(E)) = H^1(G, F_N),$$

where $F_N = \mathbf{Aut}(N)$.

The homomorphism ϱ_N is given as conjugation by elements of $\lambda(G)$.

The kernel of ϱ_N is a normal subgroup of G given as

$$G_0(N) = \{g \in G : \lambda(g)\eta\lambda(g^{-1}) = \eta, \forall \eta \in N\}.$$

The quotient group $G/G_0(N)$ is isomorphic to a subgroup U_N of F_N , which is the image of ϱ_N .

Let $E_0(N) = E^{G_0(N)}$. Then $E_0(N)$ is Galois extension of K with group U_N . So the Hopf algebra form H_N of $K[N]$ determines the data $U_N, E_0(N)/K$.

Using the Pareigis induction map

$$A(U_N, \cdot) : \mathcal{G}al(K, U_N) \rightarrow \mathcal{G}al(K, F_N),$$

we compute $A(U_N, E_0(N)/K)$, which is an F_N -Galois extension of K .

Theorem 8 (Kohl and U).

Let E/K be a Galois extension with group G and let (H_N, \cdot_N) be a Hopf-Galois structure on E/K of type N . Let

$$\Theta_N : \mathcal{G}al(K, F_N) \rightarrow \mathcal{F}orm(K[N])$$

be the Haggemüller-Pareigis bijection. Then $A(U_N, E_0(N)/K)$ is the preimage of H_N under Θ_N , i.e., $\Theta_N(A(U_N, E_0(N)/K)) = H_N$.

Proof. See [Theorem 4.3, KU25].



Example 9. Let L/\mathbb{Q} be the splitting field of the irreducible quartic $p(x) = x^4 - 10x^2 + 1$, that is, $L = \mathbb{Q}(\sqrt{2}, \sqrt{3}) = \mathbb{Q}(\sqrt{2} + \sqrt{3})$.

Then L/\mathbb{Q} is Galois with group $C_2 \times C_2 = \{1, \sigma, \tau, \tau\sigma\}$, with Galois action

$$\sigma(\sqrt{2} + \sqrt{3}) = \sqrt{2} - \sqrt{3}, \quad \tau(\sqrt{2} + \sqrt{3}) = -\sqrt{2} + \sqrt{3}.$$

By [By96], L/\mathbb{Q} admits 3 Hopf-Galois structures of type C_4 .

They correspond to regular subgroups N_1, N_2, N_3 of $\text{Perm}(G)$ normalized by $\lambda(G)$.

We first consider N_1 . As one can check

$$\begin{aligned} G_0(N_1) &= \{g \in C_2 \times C_2 : \lambda(g)\eta\lambda(g^{-1}) = \eta, \forall \eta \in N_1\} \\ &= \{1, \sigma\}. \end{aligned}$$

The fixed field of $\{1, \sigma\}$ is $\mathbb{Q}(\sqrt{2})$, and so,

$$E_0(N_1) = \mathbb{Q}(\sqrt{2}).$$

We have

$$U_{N_1} \cong (C_2 \times C_2)/G_0(N_1) = \{1, a\}, \quad a^2 = 1,$$

which is the Galois group of $E_0(N_1) = \mathbb{Q}(\sqrt{2})$.

We have $F_{N_1} = \text{Aut}(N_1) = C_2$. So in this case, the subgroup $U_{N_1} = \{1, a\}$ of F_{N_1} is all of F_{N_1} , $U_{N_1} = F_{N_1}$.

The induced F_{N_1} -Galois extension of \mathbb{Q} is

$$A(\{1, a\}, \mathbb{Q}(\sqrt{2})/\mathbb{Q}) = \mathbb{Q}(\sqrt{2}),$$

which is the preimage of H_{N_1} under the Haggemüller-Pareigis bijection

$$\Theta_{N_1} : \text{Gal}(\mathbb{Q}, F_{N_1}) \rightarrow \text{Form}(\mathbb{Q}[N_1]).$$

We next consider N_2 . Let H_{N_2} be the \mathbb{Q} -Hopf algebra attached to the Hopf-Galois structure corresponding to N_2 .

Now, H_{N_2} is a form of $\mathbb{Q}[N_2]$, but since N_1 and N_2 are of the same type (C_4), H_{N_2} is also a form of $\mathbb{Q}[N_1]$, through the isomorphism

$$\psi : N_2 \rightarrow N_1.$$

Let $F_{N_2} = \text{Aut}(N_2)$. Then there is an induced isomorphism

$$\hat{\psi} : F_{N_2} \rightarrow F_{N_1}$$

defined as $\hat{\psi}(f)(\eta) = (\psi f \psi^{-1})(\eta)$, for $f \in F_{N_2}$, $\eta \in N_1$.

The composition

$$\hat{\psi}_{\varrho_{N_2}} : G \rightarrow F_{N_2} \rightarrow F_{N_1}$$

is the homomorphism in $H^1(G, F_{N_1})$ corresponding to the form H_{N_2} of $\mathbb{Q}[N_1]$.

The kernel of $\hat{\psi}_{\varrho_{N_2}}$ is

$$\begin{aligned} G_0(N_2) &= \{g \in C_2 \times C_2 : \lambda(g)\eta\lambda(g^{-1}) = \eta, \forall \eta \in N_2\} \\ &= \{1, \tau\}. \end{aligned}$$

The fixed field of $\{1, \tau\}$ is $\mathbb{Q}(\sqrt{3})$, and so,

$$E_0(N_2) = \mathbb{Q}(\sqrt{3}).$$

We have

$$U_{N_2} \cong (C_2 \times C_2)/G_0(N_2) = \{1, b\}, \quad b^2 = 1$$

which is the Galois group of $E_0(N_2) = \mathbb{Q}(\sqrt{3})$, as a subgroup of F_{N_2} .

As a subgroup of F_{N_1} , the Galois group of $\mathbb{Q}(\sqrt{3})$ is

$$\hat{\psi}(U_{N_2}) = F_{N_1} = U_{N_1} = \{1, a\}.$$

Note: U_{N_1} acts on $\mathbb{Q}(\sqrt{3})$ through $\hat{\psi}^{-1}$, i.e.,

$$a(\sqrt{3}) = \hat{\psi}^{-1}(a)(\sqrt{3}) = b(\sqrt{3}) = -\sqrt{3},$$

as expected.

With the data $\{1, a\}$, $\mathbb{Q}(\sqrt{3})/\mathbb{Q}$, the induced F_{N_1} -Galois extension of \mathbb{Q} is

$$A(\{1, a\}, \mathbb{Q}(\sqrt{3})/\mathbb{Q}) = \mathbb{Q}(\sqrt{3}),$$

and we have

$$\Theta_{N_1}(\mathbb{Q}(\sqrt{3})) = H_{N_2}.$$

Finally, we consider N_3 . Let H_{N_3} be the \mathbb{Q} -Hopf algebra attached to the Hopf-Galois structure corresponding to N_3 .

Now, H_{N_3} is a form of $\mathbb{Q}[N_3]$, but since N_1 and N_3 are of the same type (C_4), H_{N_3} is also a form of $\mathbb{Q}[N_1]$, through an isomorphism

$$\psi : N_3 \rightarrow N_1.$$

Let $F_{N_3} = \text{Aut}(N_3)$. Then, as before, there is an induced isomorphism

$$\hat{\psi} : F_{N_3} \rightarrow F_{N_1}.$$

The composition

$$\hat{\psi}_{\varrho_{N_3}} : G \rightarrow F_{N_3} \rightarrow F_{N_1}$$

is the homomorphism in $H^1(G, F_{N_1})$ corresponding to the form H_{N_3} of $\mathbb{Q}[N_1]$.

The kernel of $\hat{\psi}_{\varrho_{N_3}}$ is

$$\begin{aligned} G_0(N_3) &= \{g \in C_2 \times C_2 : \lambda(g)\eta\lambda(g^{-1}) = \eta, \forall \eta \in N_3\} \\ &= \{1, \tau\sigma\}. \end{aligned}$$

The fixed field of $\{1, \tau\sigma\}$ is $\mathbb{Q}(\sqrt{6})$, and so,

$$E_0(N_3) = \mathbb{Q}(\sqrt{6}).$$

We have

$$U_{N_3} \cong (C_2 \times C_2)/G_0(N_3) = \{1, c\}, \quad c^2 = 1,$$

which is the Galois group of $E_0(N_3) = \mathbb{Q}(\sqrt{6})$, as a subgroup of F_{N_3} .

As a subgroup of F_{N_1} , the Galois group of $\mathbb{Q}(\sqrt{6})$ is

$$\hat{\psi}(U_{N_3}) = F_{N_1} = U_{N_1} = \{1, a\}.$$

Note: U_{N_1} acts on $\mathbb{Q}(\sqrt{6})$ through $\hat{\psi}^{-1}$, i.e.,

$$a(\sqrt{6}) = \hat{\psi}^{-1}(a)(\sqrt{6}) = c(\sqrt{6}) = -\sqrt{6},$$

as expected.

With the data $\{1, a\}$, $\mathbb{Q}(\sqrt{6})/\mathbb{Q}$, the induced F_{N_1} -Galois extension of \mathbb{Q} is

$$A(\{1, a\}, \mathbb{Q}(\sqrt{6})/\mathbb{Q}) = \mathbb{Q}(\sqrt{6}),$$

and we have

$$\Theta_{N_1}(\mathbb{Q}(\sqrt{6})) = H_{N_3}.$$

From the definition of Θ_{N_1} , we obtain

$$H_{N_1} = \mathbb{Q}(\sqrt{2})[N_1]^{\{1,a\}}$$

$$H_{N_2} = \mathbb{Q}(\sqrt{3})[N_1]^{\{1,a\}}$$

$$H_{N_3} = \mathbb{Q}(\sqrt{6})[N_1]^{\{1,a\}}$$

The H_{N_i} , $1 \leq i \leq 3$, are pairwise non-isomorphic as \mathbb{Q} -Hopf algebras since they are associated to the preimages in $\mathcal{G}al(\mathbb{Q}, F_{N_1})$: $\mathbb{Q}(\sqrt{2})/\mathbb{Q}$, $\mathbb{Q}(\sqrt{3})/\mathbb{Q}$, $\mathbb{Q}(\sqrt{6})/\mathbb{Q}$, respectively, which are clearly pairwise non-isomorphic as F_{N_1} -Galois extensions of \mathbb{Q} .

□

5. Another Example

Taylor and Truman [TT19] provide some additional examples.

Let L/K be a Galois extension of fields with quaternion group $Q_8 = \{\pm 1, \pm i, \pm j, \pm k\}$. Taylor and Truman have enumerated the Hopf-Galois structures on L/K of each possible type [Table 1, TT19].

Example 10. [Lemma 3.1, TT19]. There are 2 Hopf-Galois structures of type $C_2 \times C_2 \times C_2$.

Note that $\text{Aut}(C_2 \times C_2 \times C_2) = \text{GL}_3(\mathbb{Z}_2)$, a group of order 168.

Let M_1, M_2 be the regular subgroups of $\text{Perm}(Q_8)$ normalized by $\lambda(Q_8)$. Let H_{M_1}, H_{M_2} be the associated K -Hopf algebras.

We have

$$G_0(M_1) = G_0(M_2) = \{1, -1\}$$

and

$$E_0(M_1) = E_0(M_2) = K(\alpha, \beta),$$

the unique biquadratic subfield of L/K with Galois group

$$U_{M_1} = U_{M_2} = \mathbb{Q}_8 / \{1, -1\} = C_2 \times C_2.$$

Let $\varrho_{M_1} : \mathbb{Q}_8 \rightarrow F_{M_1}$ be the homomorphism corresponding to the L -form H_{M_1} of $K[M_1]$. The kernel of ϱ is $G_0(M_1)$; its image is U_{M_1} . We have

$$A(U_{M_1}, K(\alpha, \beta)/K) = \underbrace{K(\alpha, \beta) \times K(\alpha, \beta) \times \cdots \times K(\alpha, \beta)}_{42}$$

$$\Theta_{M_1}(A(U_{M_1}, K(\alpha, \beta)/K)) = H_{M_1}.$$

Considering the other regular subgroup M_2 , there exists a Q_8 -invariant isomorphism $\psi : M_2 \rightarrow M_1$, inducing the isomorphism

$$\hat{\psi} : F_{M_2} \rightarrow F_{M_1}.$$

Since ψ is Q_8 -invariant, $\hat{\psi}$ restricted to $U_{M_2} \leq F_{M_2}$ is the identity map.

The homomorphism corresponding to the L -form H_{M_2} of $K[M_1]$ is

$$\hat{\psi} \circ \varrho_{M_2} : Q_8 \rightarrow F_{M_2} \rightarrow F_{M_1}.$$

The data for H_{M_2} is $U_{M_1}, K(\alpha, \beta)/K$, so in fact,






$$\Theta_{M_1}(A(U_{M_1}, K(\alpha, \beta)/K)) = H_{M_2}.$$





Consequently, $H_{M_1} \cong H_{M_2}$ as K -Hopf algebras, as already shown in [TT19].



Thank you!

References

-  [AG60] Auslander, M.; Goldman, O. The Brauer group of a commutative ring. *Trans. Amer. Math. Soc.* **97** no. 3 (1960), 367–409.
-  [By96] Byott, N. Uniqueness of Hopf-Galois structures for separable field extensions. *Comm. Algebra.* **24**, (1996), 3217–3228.
-  [CHR65] Chase, S. U.; Harrison, D. K.; Rosenberg, A. Galois theory and Galois cohomology of commutative rings. *Mem. Amer. Math. Soc.*, no. 52, (1965), 15–33.
-  [HP86] Hagenmüller, R.; Pareigis, B. Hopf algebra forms of the multiplicative group and other groups. *manuscripta math.* **55** no. 2 (1986), 121–136.
-  [Ha65] Harrison, D. K. Abelian extensions of commutative rings. *Mem. Amer. Math. Soc.*, no. 52, (1965), 1–14.

-  [KKTU19b] Koch, A.; Kohl, T.; Truman, P. J.; Underwood, R. Isomorphism problems for Hopf-Galois structures on separable field extensions. *J. Pure and Appl. Algebra* **223** no. 5 (2019), 2230–2245.
-  [KU25] Kohl, T.; Underwood, R. Galois extensions and Hopf-Galois structures. *New York J. Math.* **31** (2025), 238–258.
-  [Pa90] Pareigis, B. Forms of Hopf algebras and Galois theory. *Topics in algebra, Part 1 (Warsaw, 1988)*, 75–93, Banach Center Publ., **26**, Part 1, PWN, Warsaw, 1990.
-  [TT19] Taylor, S.; Truman, P. J. The Structure of Hopf algebras giving Hopf-Galois structures on quaternionic extensions. *New York J. Math.*, **25**, (2019), 219–237.